

Childhood Lead Poisoning

You are the first line of defense
Make sure to order a Lead test

A blood lead test is the only sure way to identify a child with elevated blood lead levels.

The Centers for Disease Control and Prevention recommends blood lead testing at

12 and 24 months OR at least once before age 6

It is important to accompany diagnostic tests with education for patients on ways to reduce their children's risk for lead poisoning.



What we know — Blood Lead Level (BLL) Response Actions

There is no safe level or threshold for lead exposure

Lower levels of exposure present no obvious symptoms

Exposure may be a result of:

- Deteriorating paint from home built before 1978
- Family hobbies or occupations
- Everyday products manufactured with lead

Lead exposure may cause

- Brain and nervous system damage
- Reduced IQ
- Behavioral problems
- Learning disabilities
- Slowed growth and development
- Hearing and speech disabilities
- Premature and low birth weights in pregnant women

CDC 2018

https://www.cdc.gov/nceh/lead/acclpp/actions_blls.html

BLL Test Results (µg/dL)	Venous Confirmation Retest Within	Recommended Actions Based on BLL	Venous Retest – After Recommended Actions
< 5	None	<ul style="list-style-type: none"> • Provide dietary counseling (calcium & iron) and environmental education • Follow-up blood lead monitoring at recommended intervals 	Retest according to Blood Lead Screening Plan
5 – 9	3 mo	<p>Above Actions, plus:</p> <ul style="list-style-type: none"> • Complete history and physical exam • Lab work: Iron status, hemoglobin, hematocrit • Refer to health department for environmental investigation • Environmental assessment to identify potential lead sources • Recommend lead hazard reduction in home • Neurological, behavioral, and developmental monitoring • Abdominal x-ray with bowel decontamination (if lead ingestion is suspected) 	3 months for first 2 - 4 tests 6 - 9 months after BLL are declining
10 – 19	1 - 3 mo		1 - 3 months for first 2 - 4 tests 3 - 6 months after BLL are declining
20 – 24	1 - 3 mo		1 - 3 months for first 2 - 4 tests 1 - 3 months after BLL are declining
25 – 44	2 wk - 1 mo		2 weeks - 1 month for first 2 - 4 tests 1 month after BLL are declining
45 – 59	ASAP 48 hours		<p>Above Actions, plus:</p> <ul style="list-style-type: none"> • Oral chelation therapy • Consider hospitalization if lead-safe environment cannot be assured
60 – 69	ASAP 24 hours	Every 24 hours or as medically indicated	
≥ 70	Stat as emergency test	<p>Above Actions, plus:</p> <ul style="list-style-type: none"> • Hospitalize and commence chelation therapy in conjunction with consultation from a medical toxicologist or a pediatric environmental health specialty unit 	Every 24 hours or as medically indicated

To order educational materials for your office or families contact 702-895-1040 (Southern Nevada) | 775-453-0434 (Northern Nevada) or visit NVCLPPP.org

This publication was supported by the Grant or Cooperative Agreement Number, 1 NUE2EH001366-01-00, funded by the Centers for Disease Control and Prevention. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention or the Department of Health and Human Services.